arXiv:1704.00707v1 [math.RT] 3 Apr 2017

Critical classes, Kronecker products of spin characters, and the Saxl conjecture

Christine Bessenrodt

Institute for Algebra, Number Theory and Discrete Mathematics

Leibniz Universität Hannover
Welfengarten 1, D-30167 Hannover, Germany

bessen@math.uni-hannover.de

April 3, 2017

Abstract

Highlighting the use of critical classes, we consider constituents in Kronecker products, in particular of spin characters of the double covers of the symmetric and alternating groups. We apply results from the spin case to find constituents in Kronecker products of characters of the symmetric groups. Via this tool, we make progress on the Saxl conjecture; this claims that for a triangular number n, the square of the irreducible character of the symmetric group S_n labelled by the staircase contains all irreducible characters of S_n as constituents. We find a large number of constituents in this square which were not detected by other methods. Moreover, the investigation of Kronecker products of spin characters inspires a spin variant of Saxl's conjecture.

1 Introduction

Even though tensor products of complex representations and the corresponding Kronecker products of characters of the symmetric group S_n have been studied for a long time, their decomposition is still an elusive central open problem in the field. A new benchmark for this is a conjecture by Heide, Saxl, Tiep and Zalesskii [9] which says that for any $n \neq 2, 4, 9$ there is always an irreducible character of S_n whose square contains all irreducible characters. For triangular numbers an explicit

[0]Mathematics Subject Classifications: 20C30, 05E15.
Keywords: symmetric groups, double cover groups, characters, spin characters, Kronecker products, Saxl conjecture, unimodal sequences.

candidate was suggested by Saxl in 2012; denoting the irreducible character of S_n associated to a partition λ of n by $[\lambda]$, his conjecture is the following.

Saxl's Conjecture Let $\rho_k = (k, k-1, \ldots, 2, 1)$ be the staircase partition of $n = k(k+1)/2$. Then the Kronecker square $[\rho_k]^2$ contains all irreducible characters of S_n as constituents.

This has inspired a lot of recent research. In particular, in the work of Pak, Panova and Vallejo [19] and Ikenmeyer [12] many constituents of the square $[\rho_k]^2$ have been identified, notably those to hooks and to partitions comparable to the staircase in dominance order.

Here, we will take a very different approach and show that results on the spin characters of a double cover $\widetilde{S}_n$ of the symmetric group S_n can be fruitfully applied towards this problem. Indeed, it will turn out to be useful to consider also the spin characters of the double cover $\widetilde{A}_n$ of the alternating group A_n. An important link is made as a consequence of an identity found in an earlier investigation of homogeneous Kronecker products. While in [1] it was shown that there are no nontrivial homogeneous products of S_n-characters, for the double covers $\widetilde{S}_n$ of the symmetric groups nontrivial homogeneous spin products do occur for all triangular numbers n [2]. In [3], also nontrivial homogeneous mixed products of complex characters for the double covers $\widetilde{S}_n$ were found, i.e., products of a non-faithful character of $\widetilde{S}_n$ with a spin character. This information on nontrivial homogeneous Kronecker products is crucially used here towards obtaining many constituents in the square $[\rho_k]^2$.

We give a brief overview on the following sections. In Section 2, we collect the information and notation on the irreducible characters for the symmetric and alternating groups and their double covers. Here we already draw attention to special conjugacy classes of these groups, and we prove a general lemma that uses critical properties of conjugacy classes for identifying constituents in certain Kronecker products. In Section 3 we first recall properties of the basic spin character as well as an important link between the faithful and non-faithful characters of $\widetilde{S}_n$ labelled by the staircase partition ρ_k; as an easy application we obtain all hook characters in $[\rho_k]^2$. Section 4 starts with a short argument for the criterion given in [19, "Main Lemma"], and then moves on to the main results on constituents in certain spin character products (Theorems 4.3 and 4.11). These results are then applied towards the Saxl conjecture. A powerful new criterion for constituents in $[\rho_k]^2$ is given in Corollary 4.4; its usefulness is illustrated by providing several families of constituents. For characters to 2-part partitions this also involves the almost strict unimodality of the number of k-bounded strict partitions. The final Section 5 discusses a spin variant of Saxl's conjecture, involving the "spin staircase" $(2k-1, 2k-3, \ldots, 3, 1)$. Also spin variants of the conjecture by Heide, Saxl, Tiep and Zalesskii are presented.

2 Preliminaries

We denote by $P(n)$ the set of partitions of n, i.e., weakly decreasing sequences of nonnegative integers summing to n. For a partition $\lambda \in P(n)$, $l(\lambda)$ denotes its length, i.e., the number of positive parts of λ. The set of partitions of n into odd parts only is denoted by $O(n)$, and the set of partitions of n into distinct parts is denoted by $D(n)$. We write $D^+(n)$ resp. $D^-(n)$ for the sets of partitions λ in $D(n)$ with $n - l(\lambda)$ even resp. odd; the partition λ is then also called even resp. odd.

We write S_n for the symmetric group on n letters, and $\widetilde{S}_n$ for one one of its double covers; so $\widetilde{S}_n$ is a non-split extension of S_n by a central subgroup $\langle z \rangle$ of order 2. It is wellknown that the representation theory of these double covers is 'the same' for all representation theoretical purposes. For $\lambda \in P(n)$, we write $[\lambda]$ for the corresponding irreducible character of S_n; this is identified with the corresponding non-faithful character of $\widetilde{S}_n$. When we evaluate $[\lambda]$ on an element of S_n of cycle type $\mu \in P(n)$, we simply write $[\lambda](\mu)$ for the corresponding value. For background on the representation theory of the symmetric groups, the reader is referred to [13, 14].

The spin characters of $\widetilde{S}_n$ are those that do not have z in their kernel. For an introduction to the properties of spin characters resp. for some results we will need in the sequel we refer to [10], [16], [17], [25], [27]. Below we collect some of the necessary notation and some results from [27] that are crucial in later sections. For $n \leq 3$, the irreducible $\widetilde{S}_n$-characters are lifted from S_n. Hence we will always assume that $n \geq 4$.

Let $\lambda \in P(n)$. Then the conjugacy class of elements in $\widetilde{S}_n$ projecting to cycle type λ splits into two $\widetilde{S}_n$-conjugacy classes if and only if $\lambda \in O(n) \cup D^-(n)$, otherwise it does not split. When it splits, a specific labelling $C_\lambda^\pm$ for the two $\widetilde{S}_n$-classes in C_λ is made; we leave out the details here (see [25, 27]), and just note that for a given spin character the values on the two classes differ only by a sign. Note that any spin character vanishes on the non-split classes, thus only the values on the split classes will be considered.

Let sgn denote the sign character of $\widetilde{S}_n$, inflated from the sign character of S_n. A character χ of $\widetilde{S}_n$ is called *self-associate* if sgn $\cdot \chi = \chi$, otherwise we have a pair of *associate* characters, $\chi \neq \text{sgn} \cdot \chi$.

In 1911, Schur has proved the following classification result [25], giving a complete list of irreducible complex spin characters of $\widetilde{S}_n$.
For each $\lambda \in D^+(n)$ there is a self-associate spin character $\langle \lambda \rangle$, and for each $\lambda \in \mathcal{D}^-(n)$ there is a pair of associate spin characters $\langle \lambda \rangle_+$ and $\langle \lambda \rangle_-$. When we want to consider a spin character labelled by λ, and it is not specified whether λ is in D^+ or D^-, we write $\langle \lambda \rangle_{(\pm)}$.
The spin characters labelled by $\lambda = (\ell_1, \ldots, \ell_m) \in D(n)$ take the following values

on $\sigma_\alpha \in C_\alpha^+$:

$$\begin{aligned}
\langle\lambda\rangle_+(\sigma_\alpha) &= \langle\lambda\rangle_-(\sigma_\alpha) && \text{for } \alpha \in O(n), \lambda \in D^-(n) \\
\langle\lambda\rangle_{(\pm)}(\sigma_\alpha) &= 0 && \text{for } \alpha \in D^-(n), \lambda \neq \alpha \\
\langle\lambda\rangle_+(\sigma_\lambda) &= -\langle\lambda\rangle_-(\sigma_\lambda) = i^{(n-m+1)/2}\sqrt{\tfrac{\prod_j \ell_j}{2}} && \text{for } \lambda \in D^-(n)
\end{aligned}$$

The values $\langle\lambda\rangle_{(\pm)}(\sigma_\alpha)$ for $\alpha \in O(n)$ are integers determined by a recursion rule akin to the Murnaghan-Nakayama rule, which is due to Morris [16, 17].

Furthermore, we define

$$\widehat{\langle\lambda\rangle} = \begin{cases} \langle\lambda\rangle & \text{if } \lambda \in D^+(n) \\ \langle\lambda\rangle_+ + \langle\lambda\rangle_- & \text{if } \lambda \in D^-(n) \end{cases}$$

An important role in the theory is taken by the *basic spin characters* $\langle n\rangle_{(\pm)}$. Their values on an element $\sigma \in \widetilde{S}_n$ projecting to cycle type α are determined explicitly as follows.
When n is odd,

$$\langle n\rangle(\sigma) = 2^{(l(\alpha)-1)/2} \quad \text{for } \alpha \in O(n)\,,$$

and when $n = 2k$ is even,

$$\langle n\rangle_\pm(\sigma) = \begin{cases} 2^{(l(\alpha)-2)/2} & \text{for } \alpha \in O(n) \\ \pm i^k\sqrt{k} & \text{for } \alpha = (n) \end{cases} .$$

All other values of $\langle n\rangle_{(\pm)}$ are zero.

We will also need some information about the characters of the alternating groups A_n and their double covers $\widetilde{A}_n$.

The classification of the irreducible A_n-characters is derived from that for S_n. We obtain all irreducible characters of A_n as constituents in the restriction of the characters $[\lambda]$ as follows (see [14]).
Let $\mu \in P(n)$, and let μ' be the transposed partition to μ.
When $\mu \neq \mu'$, $[\mu]\downarrow_{A_n} = [\mu']\downarrow_{A_n} = \{\mu\} = \{\mu'\} \in \mathrm{Irr}(A_n)$.
When $\mu = \mu'$, $[\mu]\downarrow_{A_n} = \{\mu\}_+ + \{\mu\}_-$; the characters $\{\mu\}_\pm$ are conjugate irreducible characters of A_n. Let $h(\mu) = (h_1, \ldots, h_k)$ be the partition of principal hook lengths $h_1, \ldots, h_k$ in μ, i.e. here, $h_j = 2(\mu_j - j) + 1$. The two characters $\{\mu\}_\pm$ differ only on the two conjugacy classes of cycle type $h(\mu)$; note that $[\mu]$ takes the value $e_\mu = (-1)^{(n-k)/2}$ on elements of this cycle type. With $\sigma^\pm_{h(\mu)}$ being appropriate representatives of the conjugacy classes of A_n of this cycle type, the character values are given as

$$\{\mu\}_+(\sigma^\pm_{h(\mu)}) = \frac{1}{2}\left(e_\mu \pm \sqrt{e_\mu \prod_{j=1}^{k} h_j}\right)$$

and similarly, with interchanged signs, for $\{\mu\}_-$. In particular, these two conjugate characters are the only irreducible characters that differ on the elements $\sigma^\pm_{h(\mu)}$.

One also obtains the classification of irreducible spin characters of $\widetilde{A}_n$ from that of the spin characters of $\widetilde{S}_n$ (see [10, 25, 27]).
For each $\lambda \in D^-(n)$, the restriction to $\widetilde{A}_n$ gives one irreducible spin character

$$\langle\lambda\rangle_+ \downarrow_{\widetilde{A}_n} = \langle\lambda\rangle_- \downarrow_{\widetilde{A}_n} = \langle\langle\lambda\rangle\rangle \, .$$

Dually, for each $\lambda = (\ell_1, \ldots, \ell_m) \in D^+(n)$, the restriction to $\widetilde{A}_n$ gives two conjugate irreducible spin characters

$$\langle\lambda\rangle \downarrow_{\widetilde{A}_n} = \langle\langle\lambda\rangle\rangle_+ + \langle\langle\lambda\rangle\rangle_- \, .$$

If $\sigma \in \widetilde{A}_n$ projects to cycle type λ, then for the difference of the values on σ we have (with the sign depending on the choice of associates)

$$\Delta^\lambda(\sigma) = \langle\langle\lambda\rangle\rangle_+(\sigma) - \langle\langle\lambda\rangle\rangle_-(\sigma) = \pm i^{(n-m)/2} \sqrt{\prod_{j=1}^{m} \ell_j} \, ;$$

if $\sigma \in \widetilde{A}_n$ does not project to type λ, then $\Delta^\lambda(\sigma) = 0$ [25]. Let σ project to type λ. If $\lambda \in D \cap O(n)$, and $\tau \in \widetilde{A}_n$ is $\widetilde{S}_n$-conjugate to σ but not in $\sigma^{\widetilde{A}_n}$, then $\Delta^\lambda(\sigma) = -\Delta^\lambda(\tau)$ (see [27, remark after 7.5]). Note that $\langle\lambda\rangle(\sigma) = 0$ if $\lambda \in D^+(n) \setminus (D \cap O(n))$; thus we can compute all the character values also in the case where $\lambda \in D^+(n)$.
For elements projecting to other types, the values of the two conjugate characters $\langle\langle\lambda\rangle\rangle_\pm$ are the same; in particular, the characters $\langle\langle\lambda\rangle\rangle_\pm$ vanish on classes projecting to a cycle type $\mu \neq \lambda$ that is not in $O(n)$. We emphasize that when $\lambda \in D \cap O(n)$, both of these two conjugate characters have different values on the two $\widetilde{A}_n$-classes that come from one $\widetilde{S}_n$-class projecting to cycle type λ, and they are the only irreducible spin characters of $\widetilde{A}_n$ with this property.

We also note the following special situation. When $\lambda = \lambda'$, the hook length partition $h(\lambda)$ is in $D \cap O(n)$, and then the two doubling classes of $\widetilde{S}_n$ projecting to cycle type $h(\lambda)$ each split a second time, into two classes of $\widetilde{A}_n$. As pointed out above, the spin characters $\langle\langle h(\lambda)\rangle\rangle_+, \langle\langle h(\lambda)\rangle\rangle_-$ differ on the two classes of $\widetilde{A}_n$ contained in one $\widetilde{S}_n$-class but projecting onto different A_n-classes of cycle type $h(\lambda)$. Note that also the non-faithful characters $\{\lambda\}_\pm$ differ on these classes.

The alternating groups A_n and the double covers $\widetilde{S}_n$ and $\widetilde{A}_n$ exhibit a special phenomenon that we want to highlight here.

Definition 2.1 *Let G be a finite group, $I \subset \mathrm{Irr}(G)$. A pair of conjugacy classes x^G and y^G is said to be* critical *for I if $\chi(x) \neq \chi(y)$, for $\chi \in I$, but $\chi(x) = \chi(y)$ for all $\chi \in \mathrm{Irr}(G) \setminus I$.*
We call a critical pair x^G and y^G for $I = \{\chi_1, \chi_2\} \subset \mathrm{Irr}(G)$ a pair of detecting classes *for χ_1, χ_2 if $\chi_1(x) - \chi_1(y) = \chi_2(y) - \chi_2(x)$.*

Of course, the set I above should be taken of small size to give interesting information.

Remarks and Examples 2.2 (i) If there is a pair of detecting classes for χ_1, χ_2, then $\chi_2 = \overline{\chi_1}$, or the characters are both real.

(ii) For A_n, for each symmetric partition μ of n, the pair of classes $\sigma^{\pm}_{h(\mu)}$ is a detecting pair for $\{\mu\}_{\pm}$.

(iii) For $\widetilde{S}_n$, for each $\lambda \in D^-(n)$, the pair of classes of elements projecting to cycle type λ is a detecting pair for $\langle\lambda\rangle_{\pm}$.

(iv) For $\widetilde{A}_n$, for each $\lambda \in D^+(n)$, the two pairs of classes of elements of $\widetilde{A}_n$ projecting to cycle type λ and belonging to one $\widetilde{S}_n$ class are detecting pairs for $\langle\langle\lambda\rangle\rangle_{\pm}$.

(v) While the situations above are the ones used in the later sections, it should be noted that there many more such instances, and we mention just a few examples. For $G = \mathrm{GL}(3,2)$, the pair of classes of elements of order 7 is detecting for the pair of characters of degree 3. For $G = \mathrm{PSL}(2,11)$, the pair of classes of elements of order 5 (order 11, resp.) is detecting for the pair of characters of degree 12 (degree 5, resp.). For $G = M_{11}$, the pair of classes of elements of order 8 (order 11, resp.) is detecting for the conjugate pair of characters of degree 10 (degree 16, resp.).

The reason for the notions of critical and detecting classes is the following easy but very useful result on Kronecker products. It originates with the usage of detecting classes in [4], its variations in [19, 20], and the idea to consider detecting classes for pairs of spin characters of the double cover groups; we will follow this up in later sections.

Lemma 2.3 *Let G be a finite group, $x, y \in G$. Let ψ be a character of G with $\psi(x) = \psi(y) \neq 0$.*

(1) Assume that x^G, y^G is a critical pair for $I \subset \mathrm{Irr}(G)$. Then for any $\chi \in I$, $\psi \cdot \chi$ has a constituent in I.
In particular, if ψ is irreducible and $\chi \in I$, then $\chi \cdot \sum_{\nu \in I} \nu$ contains ψ as a constituent.

(2) Assume that x^G, y^G is a detecting pair for $\chi_1, \chi_2 \in \mathrm{Irr}(G)$. Set $m_j = \langle \psi \cdot \chi_1, \chi_j \rangle$, $j = 1, 2$. Then

$$\psi(x) = m_1 - m_2 \,.$$

Furthermore,

$$\max(m_1, m_2) \geq |\psi(x)| > 0 \,.$$

Proof. The assertions in (1) are obvious (note that I is closed under complex conjugation). For the claim in (2), we compute the difference of the values of $\psi \cdot \chi_j$ on the two classes. Set $t := \chi_1(x) - \chi_1(y) = \chi_2(y) - \chi_2(x)$; note that $t \neq 0$. First we have

$$\psi\chi_1(x) - \psi\chi_1(y) = \psi(x)(\chi_1(x) - \chi_1(y)) = \psi(x)t.$$

On the other hand, $\psi\chi_1 = m_1\chi_1 + m_2\chi_2 + \sum_{\chi\neq\chi_1,\chi_2} m_\chi\chi$, and hence (as the pair x^G, y^G is detecting for χ_1, χ_2) we obtain

$$\psi\chi_1(x) - \psi\chi_1(y) = m_1(\chi_1(x) - \chi_1(y)) + m_2(\chi_2(x) - \chi_2(y)) = (m_1 - m_2)t.$$

As $t \neq 0$, we deduce $\psi(x) = m_1 - m_2$. The assertion $\max(m_1, m_2) \geq |\psi(x)|$ now follows immediately. $\diamond$

3 The basic spin character and hooks

Of central interest in the representation theory of the symmetric groups are the Kronecker coefficients $g(\lambda, \mu, \nu)$ appearing as expansion coefficients in the Kronecker products

$$[\lambda][\mu] = \sum_\nu g(\lambda, \mu, \nu)[\nu]\,.$$

Using this notation, Saxl's conjecture may be rephrased as saying the following for the staircase partition $\rho_k = (k, k-1, \ldots, 2, 1)$ of $n = k(k+1)/2$:

$$g(\rho_k, \rho_k, \lambda) > 0 \quad \text{for all partitions } \lambda \text{ of } n\,.$$

As we will show in the following, products of spin characters or mixed products of a spin character and a non-faithful character can play an important role towards finding constituents in the square $[\rho_k]^2$.

For the product of any ordinary character $[\lambda]$ of S_n with the basic spin characters $\langle n\rangle_{(\pm)}$, Stembridge has provided an efficient combinatorial formula in [27]. It was already observed in [3] that as an immediate consequence of this formula the following result is obtained.

Proposition 3.1 *We have the following spin product decomposition:*

$$\langle n\rangle \cdot \widehat{\langle n\rangle} = \sum_{j=0}^{n-1} [n-j, 1^j] =: \chi_{hook}.$$

Regev [21] showed that the sum of all hook characters has particular easy values; Taylor recently gave a different proof of this fact [28]. Here, we point out that this is a direct consequence of the proposition above and the knowledge of the values of the basic spin characters stated in Section 2:

Corollary 3.2 *Let $\sigma \in S_n$ be of cycle type α. Then*

$$\chi_{hook}(\sigma) = \begin{cases} 2^{l(\alpha)-1} & \text{if } \alpha \in O(n) \\ 0 & \text{otherwise} \end{cases}\,.$$

For our purpose of finding constituents in the square of the staircase character the following result on spin products for $\widetilde{S}_n$ turns out to be important; it was obtained in the context of classifying the homogeneous spin products [2]. Here, it provides the crucial link to the Saxl conjecture.

Proposition 3.3 *Let $n = k(k+1)/2$, $\rho_k = (k, k-1, \ldots, 2, 1)$. Then*

$$\langle n \rangle_{(\pm)} \cdot \langle \rho_k \rangle_{(\pm)} = 2^{a(k)} [\rho_k]$$

where $a(k)$ is given by

$$a(k) = \begin{cases} \frac{k-2}{2} & \text{if } k \text{ is even} \\ \frac{k-1}{2} & \text{if } k \equiv 1 \mod 4 \\ \frac{k-3}{2} & \text{if } k \equiv 3 \mod 4 \end{cases} .$$

The proposition above is the key for finding new constituents in $[\rho_k]^2$; this is due to the following observation.

Lemma 3.4 *Let $[\lambda]$ be a constituent of $\langle \rho_k \rangle_{(\pm)} \cdot \langle \rho_k \rangle_{(\pm)}$, for a suitable choice of associates. Then all constituents of $\chi_{hook} \cdot [\lambda]$ are constituents of $[\rho_k]^2$. In particular, $[\lambda]$ is a constituent of $[\rho_k]^2$.*

Proof. For any choice of associates we have

$$\langle n \rangle_{(\pm)} \langle n \rangle_{(\pm)} \cdot \langle \rho_k \rangle_{(\pm)} \cdot \langle \rho_k \rangle_{(\pm)} = 2^{2a(k)} [\rho_k]^2 . \tag{1}$$

Now the assertion follows immediately from Proposition 3.1 $\diamond$

As a contribution towards the Saxl conjecture we immediately obtain the following result, which was proved by very different methods in [12] and a weaker asymptotic version in [19]:

Corollary 3.5 *All hook characters $[n-j, 1^j]$ are constituents in $[\rho_k]^2$.*

Proof. For any $\mu \in D(n)$, $\overline{\langle \mu \rangle_{(\pm)}}$ is one of the characters $\langle \mu \rangle_{(\pm)}$, so the product $\langle \mu \rangle_{(\pm)} \cdot \langle \mu \rangle_{(\pm)}$ contains $[n]$ for a suitable choice of associates. Hence the assertion follows immediately from Lemma 3.4. $\diamond$

Remark 3.6 Applying Lemma 3.4 with other constituents $[\lambda]$ than the trivial character, gives further near-hook constituents; we will see more such constituents in the next section. For an example at this point, using the constituents $[n-3, 3]$ or $[n-3, 1^3]$ of $\langle \rho_k \rangle_{(\pm)} \cdot \langle \rho_k \rangle_{(\pm)}$, for a suitable choice of associates, obtained in [3, Theorem 3.5], one finds as constituents in $[\rho_k]^2$ characters to double-hooks with the smaller principal hook being of size at most 3.

4 Constituents in spin products and the Saxl conjecture

As mentioned earlier, the common theme in the character theories of the alternating groups and the double covers of the symmetric and alternating groups that is crucial in the applications here is the existence of critical or detecting classes.

For example, by using the characters of the alternating groups, it was shown in [4] that for a symmetric partition λ, the character $[\lambda]$ is always a constituent in its own square $[\lambda]^2$. The idea was to use the pair of conjugacy classes in the alternating groups that detect the characters $\{\lambda\}_\pm$, namely the ones of cycle type $h(\lambda)$. This was taken up by Pak, Panova and Vallejo in [19] to provide a criterion for constituents in $[\lambda]^2$; we give a very short argument here.

Lemma 4.1 *[19, "Main Lemma"] Let λ be a symmetric partition. Let μ be a partition with $[\mu](h(\lambda)) \neq 0$. Then $[\mu]$ is a constituent of $[\lambda]^2$.*

Proof. The restriction $\chi = [\mu]\downarrow_{A_n}$ takes the same (non-zero) values on the two classes of cycle type $h(\lambda)$, which are critical for the pair of characters $\{\lambda\}_\pm$. Hence by Lemma 2.3 the character $\chi \cdot \{\lambda\}_+$ contains one of $\{\lambda\}_\pm$, and thus the covering product $[\mu] \cdot [\lambda]$ contains $[\lambda]$, i.e., $g(\lambda, \lambda, \mu) > 0$, as required. $\diamond$

Towards the Saxl conjecture this immediately gives:

Corollary 4.2 *[19] Let μ be a partition such that $[\mu](h(\rho_k)) \neq 0$. Then $[\mu]$ is a constituent of $[\rho_k]^2$.*

Unfortunately, many irreducible characters vanish on the class of cycle type $h(\rho_k)$; this was already analysed in [19]. It turns out that the idea of critical detecting classes is more powerful in the context of spin characters.

Theorem 4.3 *Let $\lambda \in D(n)$, $n \geq 4$. Let μ be a partition such that $[\mu](\lambda) \neq 0$. Then $[\mu]$ is a constituent of $\langle\lambda\rangle_{(\pm)}\widehat{\langle\lambda\rangle}$, i.e., of $\langle\lambda\rangle_{(\pm)} \cdot \langle\lambda\rangle_{(\pm)}$ for a suitable choice of associates.*

Proof. For $\lambda \in D^-(n)$, this follows immediately from Lemma 2.3(1), as the two $\widetilde{S}_n$-classes projecting to cycle type λ are a critical pair for the pair of characters $\langle\lambda\rangle_\pm$.

For $\lambda \in D^+(n)$, we use spin characters of the double covers of the alternating groups. First assume $\lambda \in D^+(n) \setminus (D \cap O(n))$. Then the two classes of $\widetilde{A}_n$ projecting to cycle type λ are critical for the two spin characters of $\widetilde{A}_n$ labelled by λ. Hence one of $\langle\langle\lambda\rangle\rangle_\pm$ is a constituent of $[\mu]\downarrow_{\widetilde{A}_n} \cdot\langle\langle\lambda\rangle\rangle_\pm$; thus $\langle\lambda\rangle$ is a constituent of $[\mu] \cdot \langle\lambda\rangle$, and the claim follows. Now take $\lambda \in D \cap O(n)$; note that we then have four $\widetilde{A}_n$-classes projecting to type λ. We consider non-conjugate elements $\sigma_1, \sigma_2 \in \widetilde{A}_n$ that belong to the same $\widetilde{S}_n$-class, projecting to cycle type λ. Recall that $\langle\langle\lambda\rangle\rangle_\pm$ are the only irreducible spin characters that differ on σ_1, σ_2. Now $[\mu]\downarrow_{\widetilde{A}_n} \cdot\langle\langle\lambda\rangle\rangle_\pm$ is a

spin character with different values on σ_1, σ_2, hence it must have one of $\langle\langle\lambda\rangle\rangle_\pm$ as a constituent, implying the claim as before. $\diamond$

This leads to a powerful criterion in the context of the Saxl conjecture:

Corollary 4.4 *Let μ be a partition such that $[\mu](\rho_k) \neq 0$. Then $[\mu]$ is a constituent of $[\rho_k]^2$.*

Proof. The assertion follows by Lemma 3.4 and Theorem 4.3. $\diamond$

Remark 4.5 Using non-vanishing on the class ρ_k produces many more constituents than non-vanishing on the class $h(\rho_k)$! Fortunately, the two tests may be combined to give even more constituents. For an illustration of this, here are the numerical values up to $k = 11$:

k	n	$p(n)$	non-zero on $h(\rho_k)$	non-zero on ρ_k	non-zero on ρ_k or on $h(\rho_k)$	%
2	3	3	3	2	3	100
3	6	11	5	6	9	81.8
4	10	42	21	24	33	78.6
5	15	176	45	114	131	74.4
6	21	792	231	524	607	76.6
7	28	3718	573	2408	2623	70.5
8	36	17977	3321	12734	13567	75.5
9	45	89134	9321	67462	69692	78.2
10	55	451276	59091	370590	381375	84.5
11	66	2323520	183989	2036486	2060003	88.7

The non-vanishing on ρ_k also seems to be a good criterion when compared with Ikenmeyer's criterion which gives the constituents to partitions comparable to ρ_k in dominance order [12]; in the region above, the corresponding percentage is decreasing, and already below 50% at $k = 9$.

We want to illustrate the usefulness of the criterion given above by showing that in particular all characters to 2-part partitions are constituents of $[\rho_k]^2$. Note that by the criterion which uses the value on the class to $h(\rho_k)$, asymptotically, these constituents are found in [19, Corollary 6.4]; on the other hand, the constituents to 2-part partitions are also obtained by Ikenmeyer's result [12].
In fact, as we will see below, by the approach via spin characters many further constituents of $[\rho_k]^2$ are obtained, namely the constituents in products of hook characters and characters to 2-part partitions.

We consider a character to a 2-part partition $[n-j,j]$, with $1 \le j \le n/2$. By the Littlewood-Richardson rule, we have

$$[n-j,j] = [n-j] \circ [j] - [n-j+1] \circ [j-1],$$

where on the right hand side the outer product of two characters is denoted by $\circ$.

For $k \le n$, we set

$$d_k(n) = |\{\lambda = (\ell_1, \dots) \in D(n) \mid \ell_1 \le k\}| \,.$$

First we want to apply the criterion given in Theorem 4.3, that is, we have to evaluate $[n-j,j]$ at ρ_k, for $n = k(k+1)/2$. By the above, we obtain the value

$$[n-j,j](\rho_k) = d_k(j) - d_k(j-1) \,.$$

The partition numbers $d_k(m)$, $0 \le m \le n = k(k+1)/2$, are easily seen to form a symmetric sequence. They are also the coefficients in the expansion

$$\prod_{i=1}^{k}(1+x^i) = \sum_{m=0}^{n} d_k(m)x^m \,.$$

This polynomial is known to be unimodal, by quite different and intricate proofs due to Hughes [11], and Odlyzko and Richmond [18]; see also Stanley's article [26] for more on this and related unimodal sequences. Here, based on a result by Odlyzko and Richmond [18], we find that the sequence is in fact *almost strictly* unimodal; the few exceptions for small k, and at the very beginning and end (and an instance in the middle when the sequence has even length) can be described explicitly.

Theorem 4.6 *For $k \in \mathbb{N}$, the following holds for the partition numbers $d_k(m)$:*

(i) For $k \ge 1$, $d_k(0) = d_k(1) = 1$.

(ii) For $k \ge 2$, $d_k(2) = 1$; $d_2(3) = 1$, $d_3(4) = 1$.

(iii) For $k \ge 4$, $d_k(3) = d_k(4) = 2$.

(iv) For $k \in \{4, \dots, 11\}$, further equalities $d_k(m-1) = d_k(m)$ hold for the following values of $m \le k(k+1)/4$:

k	4	5	6	7	8	9	10	11
m	5	6, 7	7, 8, 10	8, 11, 13, 14	16, 17	19, 22	26	32

(v) For $k \ge 12$, $d_k(m-1) < d_k(m)$ for $5 \le m \le k(k+1)/4$.

Up to equalities deduced from the above by the symmetry of the sequence (including an equality in the middle when $k \equiv 1$ or $2 \mod 4$), all equalities in the partition sequences $(d_k(m))_{0 \le m \le k(k+1)/2}$, $k \in \mathbb{N}$, are described above.

Proof. The equalities given in (i)-(iii) above are easily seen, and the exceptional ones in (iv) are obtained by computation. Up to the relations coming from the symmetry of the sequence, it only remains to discuss (v) to cover all relations between $d_k(m)$ and $d_k(m-1)$.

Considering strict partitions of m with largest part being smaller than k or equal to k, respectively, gives the recursion

$$d_k(m) = d_{k-1}(m) + d_{k-1}(m-k)\,.$$

This will now be applied to show in an induction on k, that for $k \ge 12$ the sequence $d_k(m)$ is strictly increasing in the range $4 \le m \le k(k+1)/4$.

For $12 \le k < 60$, this claim holds by direct computation. So we assume now that $k \ge 60$. Then, using unimodality of the sequences and induction, for $m \le k(k-1)/4$ we obtain

$$d_k(m) - d_k(m-1) = d_{k-1}(m) - d_{k-1}(m-1) + d_{k-1}(m-k) - d_{k-1}(m-1-k) > 0\,.$$

For the range $k(k-1)/4 \le m \le k(k+1)/4$, the inequality $d_k(m-1) < d_k(m)$ is provided by [18, Theorem 4]. Hence, from $m = 4$ onwards, the d_k-sequence increases up to the middle; for $k \equiv 1, 2 \mod 4$, there is an equality in the middle by symmetry. $\diamond$

From the discussion before the theorem we now deduce immediately:

Corollary 4.7 *Let $k \in \mathbb{N}$, $n = k(k+1)/2$. Then $[n-j, j](\rho_k) > 0$ for $0 \le j \le n/2$, unless j is one of the exceptional values given in Theorem 4.6 where $d_k(j-1) = d_k(j)$.*

Our criterion in Theorem 4.3 now implies:

Corollary 4.8 *Let $k \in \mathbb{N}$, $n = k(k+1)/2$, $0 \le j \le n/2$ such that j is not one of the exceptional values in Theorem 4.6. Then $[n-j, j]$ is a constituent of $\langle \rho_k \rangle_{(\pm)} \cdot \langle \rho_k \rangle_{(\pm)}$ for a suitable choice of associates.*

Towards the Saxl conjecture we deduce:

Corollary 4.9 *Let $k \in \mathbb{N}$, $n = k(k+1)/2$.*

(1) Assume $0 \le j \le n/2$ and j is not one of the exceptional values in Theorem 4.6. Then all constituents of $\chi_{hook} \cdot [n-j, j]$ are constituents of $[\rho_k]^2$.

(2) All characters $[n-j, j]$, $0 \le j \le n/2$, are constituents of $[\rho_k]^2$.

Proof. The first assertion follows immediately using Lemma 3.4. For the second assertion, we only have to check the characters $[n-j,j]$ for the few exceptional values of j where $[n-j,j](\rho_k)=0$. For the cases $j=1,k\geq 2$, or $j=2,k\geq 3$, or $j=4,k\geq 4$, the assertion $g(\rho_k,\rho_k,(n-j,j))>0$ is already known by work of Saxl [24]. So it only remains to check the few exceptional cases for $k\leq 11$, which is easily done by computation. $\diamond$

Remark 4.10 Products of hook characters and characters to 2-part partitions have been studied by Remmel [22] and Rosas [23]. The extra constituents that we obtain from Corollary 4.9 are all labelled by double hooks; in fact, every character to a double hook appears in a product of the form $\chi_{\rm hook}\cdot[n-j,j]$, where there is some flexibility in the choice of j [22, Theorem 2.2]. Avoiding the few exceptional values for j in Theorem 4.6 and showing $g(\rho_k,\rho_k,\lambda)>0$ for small k by computation (or one of the available criteria) then gives all double hook characters $[\lambda]$ in $[\rho_k]^2$.

Already in [4], for symmetric λ the result on the multiplicity $g(\lambda,\lambda,\lambda)$ was made more precise; from the calculation of the scalar products on the level of the alternating groups a congruence mod 4 was found for the Kronecker coefficient. Indeed, a similar calculation was used to find the value $|[\mu](h(\lambda))|$ as a lower bound for the Kronecker coefficient $g(\lambda,\lambda,\mu)$ in [20]; we have seen a general version of this in Lemma 2.3.

Also in the spin case we can make the result in Theorem 4.3 more precise.

Theorem 4.11 *Let $\lambda\in D(n)$, $n\geq 4$. Let μ be a partition such that $[\mu](\lambda)\neq 0$.*

(1) Let $\lambda\in D^-(n)$. Set $m_\pm=\langle\langle\lambda\rangle_\pm,[\mu]\cdot\langle\lambda\rangle_+\rangle$. Then

$$m_+-m_-=[\mu](\lambda)\,.$$

In particular, $0<|[\mu](\lambda)|\leq\max(m_+,m_-)\leq\langle[\mu],\langle\lambda\rangle_\pm\cdot\widehat{\langle\lambda\rangle}\rangle$.

(2) Let $\lambda\in D^+(n)$. Set $m_\pm=\langle\langle\langle\lambda\rangle\rangle_\pm,[\mu]\downarrow_{\widetilde{A}_n}\cdot\langle\langle\lambda\rangle\rangle_+\rangle$. Then

$$m_+-m_-=[\mu](\lambda)$$

and

$$m_++m_-=2m_-+[\mu](\lambda)=\langle[\mu]\langle\lambda\rangle,\langle\lambda\rangle\rangle\,.$$

In particular,

$$0<|[\mu](\lambda)|\leq\max(m_+,m_-)\leq\langle[\mu]\langle\lambda\rangle,\langle\lambda\rangle\rangle$$

and

$$\langle[\mu]\langle\lambda\rangle,\langle\lambda\rangle\rangle\equiv[\mu](\lambda)\mod 2\,.$$

Proof. The D^- case is an immediate consequence of Lemma 2.3.
For the D^+ case, we compute the difference of the values of $[\mu]\downarrow_{\widetilde{A}_n}\cdot\langle\langle\lambda\rangle\rangle_+$ on the two $\widetilde{A}_n$-classes contained in one $\widetilde{S}_n$ class projecting to cycle type λ, similarly as in the proof of Lemma 2.3. Note that this character is a linear combination of irreducible spin characters, and restricted to the spin characters, the two $\widetilde{A}_n$-classes are critical. This gives the first assertion. Observing that $\langle\langle\langle\lambda\rangle\rangle_\pm,[\mu]\downarrow_{\widetilde{A}_n}\cdot\langle\langle\lambda\rangle\rangle_-\rangle = m_\mp$ then gives the second assertion.

In both cases the additional claims are an immediate consequence. $\diamond$

Remark 4.12 There is a different way to obtain a character $[\mu]$ as a constituent in $[\rho_k]^2$ by using its character value on the class of cycle type ρ_k. As the criterion to be described now is much weaker, we only provide the main arguments without going into the necessary background in detail; again, it is based on using a special detection property of the class of cycle type ρ_k.

By the results in [5] and [6], each partition $\alpha \in O(n)$ is special for the spin character(s) $\langle\beta\rangle_{(\pm)}$ to its Glaisher correspondent $\beta \in D(n)$, with respect to the condition that the 2-power in the spin character value of $\langle\beta\rangle_{(\pm)}$ on elements projecting to cycle type α is the smallest (among the 2-powers in spin character values on this class). The spin character(s) to $\beta = \rho_k$ are indeed unique (up to associates) with this property on their special class, to the Glaisher correspondent α of ρ_k. Thus, when we multiply $\langle\rho_k\rangle_{(\pm)}$ with a character $[\mu]$ that has odd value on α, the product has to contain $\langle\rho_k\rangle_{(\pm)}$ as a constituent. Now it follows from a general character-theoretic fact [7, (6.4)] that the values $[\mu](\rho_k)$ and $[\mu](\alpha)$ are congruent modulo 2. Hence, whenever $[\mu](\rho_k)$ is odd, we deduce that $[\mu]$ is a constituent of $\langle\rho_k\rangle_{(\pm)}\widehat{\langle\rho_k\rangle}$, and hence also of $[\rho_k]^2$.

5 A spin variant of Saxl's conjecture

The product of two spin characters for the double cover groups decomposes into non-faithful irreducible characters, while a mixed product of a spin character and a non-faithful character decomposes into irreducible spin characters. Thus in a variant of Saxl's conjecture for the double cover groups we cannot expect to find an irreducible character whose square contains all irreducible characters but have to be more modest.

We start with a result obtained in the context of classifying homogeneous mixed products; it is a special product appearing in [3, Theorem 3.2]:

Proposition 5.1 *For $n = k^2 \geq 4$ we have*

$$\langle n\rangle_{(\pm)}\cdot[k^k] = 2^{\lfloor\frac{k-1}{2}\rfloor}\langle 2k-1, 2k-3, \ldots, 3, 1\rangle\,.$$

We denote by $\tau_k = (2k-1, 2k-3, \ldots, 3, 1) \in D(k^2)$ the "spin staircase" of length k; note that $\tau_k = h((k^k))$.

Arguing similarly as for Lemma 3.4 we immediately deduce from Proposition 5.1:

Lemma 5.2 *Let $[\lambda]$ be a constituent of $[k^k]^2$. Then all constituents of $\chi_{hook} \cdot [\lambda]$ are constituents of $\langle\tau_k\rangle^2$.*

In particular, the square $\langle\tau_k\rangle^2$ contains $[\lambda]$ and all hook characters $[n-j, 1^j]$, $j \in \{0, 1, \ldots, n-1\}$.

Using Lemma 4.1, we deduce from Lemma 5.2:

Corollary 5.3 *Let $n = k^2$. Then the square $\langle\tau_k\rangle^2$ contains all characters $[\mu]$ such that $[\mu](\tau_k) \neq 0$.*

Remark 5.4 In fact, for $2 \leq k \leq 5$, $\langle\tau_k\rangle^2$ contains all characters $[\mu]$, $\mu \in P(k^2)$. So as a spin variant of Saxl's conjecture we may ask whether this holds for all $k \geq 2$.

From the criterion in Corollary 5.3 above, already many constituents are obtained; here are the numerical values up to $k = 8$:

k	n	$p(n)$	non-zero on τ_k	%
1	1	1	1	100
2	4	5	3	60.0
3	9	30	15	50.0
4	16	231	93	40.3
5	25	1958	755	38.6
6	36	17977	7185	40.0
7	49	173525	75430	43.5
8	64	851522	1741630	48.9

Computations with GAP [8] led to the following conjecture, adding to the conjectures made by Heide, Saxl, Tiep and Zalesskii [9] on character squares; note that for $\lambda \in D^-(n)$, always one of $[n]$ or $[1^n]$ is missing from the square $\langle\lambda\rangle^2$.

Conjecture 5.5 *For any $n \geq 4$, $n \neq 5$, there is a spin character $\langle\lambda\rangle$, $\lambda \in D^+(n)$, whose square $\langle\lambda\rangle^2$ contains all $[\mu]$, $\mu \in P(n)$.*

It was also conjectured in [9] that for $n \geq 4$ there is always an irreducible character of the alternating group A_n whose square contains all irreducible A_n characters; in fact, a quick check with GAP up to $n = 25$ shows that for growing n a large percentage of irreducible A_n characters has this property. For the double cover groups $\widetilde{A}_n$, data computed by GAP up to $n = 25$ provides evidence for the following conjecture; again, for growing n, it seems that a large percentage of irreducible $\widetilde{A}_n$ spin characters has the property considered here.

Conjecture 5.6 *For any $n \geq 5$ there is a spin character $\chi \in \mathrm{Irr}(\widetilde{A}_n)$ whose square χ^2 contains all non-faithful $\psi \in \mathrm{Irr}(\widetilde{A}_n)$.*

References

[1] C. Bessenrodt, A. Kleshchev, On Kronecker products of complex representations of the symmetric and alternating groups. *Pacific J. Math.* 190 (1999) 201–223.

[2] C. Bessenrodt, A. Kleshchev, On Kronecker products of spin characters of the double covers of the symmetric groups. *Pacific J. Math.* 198 (2001) 295–305.

[3] C. Bessenrodt, On mixed products of complex characters of the double covers of the symmetric groups. *Pacific J. Math.* 199 (2001) 257–268.

[4] C. Bessenrodt, C. Behns, On the Durfee size of Kronecker products of characters of the symmetric groups and their double covers. *J. Algebra* 280 (2004) 132–144.

[5] C. Bessenrodt, J.B. Olsson, Spin representations and powers of 2. *Algebras Rep. Theory* 3 (2000) 289–300.

[6] C. Bessenrodt, J.B. Olsson, Spin representations, powers of 2 and the Glaisher map. *Algebras Rep. Theory* 8 (2005) 1–10.

[7] W. Feit, *Characters of finite groups*. W. A. Benjamin, Inc., New York 1967.

[8] The GAP Group, *GAP – Groups, Algorithms, and Programming, Version 4.7.4*; 2014. (\protect\vrule width0pt\protect\href{http://www.gap-system.org}{http://www.gap-syste

[9] G. Heide, J. Saxl, P. Tiep, A. Zalesskii, Conjugacy action, induced representations and the Steinberg square for simple groups of Lie type. *Proc. London Math. Soc.* 106 (2013) 908–930.

[10] P. Hoffman, J.F. Humphreys, *Projective representations of the symmetric groups.* Oxford, 1992.

[11] J.W.B. Hughes, Lie algebraic proof of some theorems on partitions. In: Number Theory and Algebra (H. Zassenhaus, ed.), 135–155, Academic Press (1977).

[12] C. Ikenmeyer, The Saxl conjecture and the dominance order. *Discrete Math.* 338 (2015) 1970–1975.

[13] G. James, *The representation theory of the symmetric groups,* Springer Lecture Notes Math. 682 (1978).

[14] G. James, A. Kerber, *The representation theory of the symmetric group,* Addison-Wesley, London, 1981.

[15] P.B. Kleidman, D.B. Wales, The projective characters of the symmetric groups that remain irreducible on subgroups, *J. Algebra* 138 (1991) 440–478.

[16] A.O. Morris, The spin representation of the symmetric group. *Proc. London Math. Soc.* (3) 12 (1962) 55–76.

[17] A.O. Morris, The spin representation of the symmetric group. *Canad. J. Math.* 17 (1965) 543–549.

[18] A.M. Odlyzko, L.B. Richmond, On the unimodality of some partition polynomials. *Europ. J. Comb.* 3 (1982) 69–84.

[19] I. Pak, G. Panova, E. Vallejo, Kronecker products, characters, partitions, and the tensor square conjectures. *Advances Math.* 288 (2013) 702–731.

[20] I. Pak, G. Panova, Bounds on certain classes of Kronecker and q-binomial coefficients, *J. Comb. Theory A* 147 (2017) 1–17.

[21] Amitai Regev, Lie superalgebras and some characters of S_n. *Israel J. Math.* 195 (2013) 31–35.

[22] J. Remmel, Formulas for the expansion of the Kronecker products $S_{(m,n)} \otimes S_{(1^{p-r},r)}$ and $S_{(1^k 2^l)} \otimes S_{(1^{p-r},r)}$. *Discrete Math.* 99 (1992) 265–287.

[23] M.H. Rosas, The Kronecker product of Schur functions indexed by 2-row shapes or hook shapes. *J. Algebraic Combin.* 14 (2001) 153–173.

[24] J. Saxl, The complex characters of the symmetric groups that remain irreducible in subgroups. *J. Algebra* 111 (1987) 210–219.

[25] I. Schur, Über die Darstellung der symmetrischen und der alternierenden Gruppen durch gebrochen lineare Substitutionen. *J. Reine und Angew. Math.* 139 (1911) 155–250.

[26] R.P. Stanley, Log-concave and unimodal sequences in algebra, combinatorics, and geometry. In: Graph Theory and Its Applications: East and West, Ann. New York Acad. Sci., vol. 576 (1989) 500–535.

[27] J. Stembridge, Shifted tableaux and the projective representations of symmetric groups. *Advances Math.* 74 (1989) 87–134.

[28] J. Taylor, A note on skew characters of symmetric groups. arXiv:1603.03914v1